

Relation between geometric phases of entangled bi-partite systems and their subsystems

D.M. Tong¹, E. Sjöqvist^{2*}, L.C. Kwek^{1,3}, C.H. Oh^{1†}, and M. Ericsson^{2,4‡}

¹*Department of Physics, National University of Singapore, 10 Kent Ridge Crescent, Singapore 119260*

²*Department of Quantum Chemistry, Uppsala University, Box 518, Se-751 20 Uppsala, Sweden*

³*National Institute of Education, Nanyang Technological University, 1 Nanyang Walk, Singapore 639798*

⁴*Department of Physics, University of Illinois at Urbana-Champaign,
1110 West Green Street, Urbana, IL 61801-3080, USA*

(Dated: February 1, 2008)

This paper focuses on the geometric phase of entangled states of bi-partite systems under bi-local unitary evolution. We investigate the relation between the geometric phase of the system and those of the subsystems. It is shown that (1) the geometric phase of cyclic entangled states with non-degenerate eigenvalues can always be decomposed into a sum of weighted non-modular pure state phases pertaining to the separable components of the Schmidt decomposition, though the same cannot be said in the non-cyclic case, and (2) the geometric phase of the mixed state of one subsystem is generally different from that of the entangled state even by keeping the other subsystem fixed, but the two phases are the same when the evolution operator satisfies conditions where each component in the Schmidt decomposition is parallel transported.

PACS numbers: 03.65.Vf, 03.67.Lx

I. INTRODUCTION

In 1984, Berry [1] proposed in a seminal paper that a quantum system in a pure state undergoing adiabatic cyclic evolution acquires a geometric phase. This discovery has prompted a myriad of activities on various aspects of geometric phase in many areas of physics ranging from optical fibers to anyons. Simon [2] subsequently recast the mathematical formalism of Berry's phase within the language of differential geometry and fiber bundles. While it is possible to consider Berry's phase under adiabatic evolution, the extension to non-adiabatic evolution is usually non-trivial. The general formalism for the non-adiabatic extension was formulated by Aharonov and Anandan [3, 4]. Samuel and Bhandari [5] further generalized the geometric phase by extending it to non-cyclic evolution and sequential measurements. Further relaxation on the adiabatic, unitary, and cyclic properties of the evolution have since been carried out [6, 7, 8].

The concept of geometric phase of mixed states has also been developed. Uhlmann [9] was probably first to address this issue within the mathematical context of purification. Sjöqvist *et al.* [10] have introduced a new formalism that defines the mixed state geometric phase within the experimental context of quantum interferometry. As pointed out by Slater [11], these two approaches are not equivalent, and Ericsson *et al.* [12] have recently shown that the parallel transport conditions used in the two approaches lead to generically distinct phase holonomy effects for entangled systems undergoing certain local unitary transformations. Singh *et al.* [13] have given a kinematic formalism of the mixed state geometric phase with non-degenerate eigenvalues and generalized the analysis of Ref. [10] to degenerate states. Extension of Ref. [10] to the off-diagonal case [14] and completely positive maps [15] have also been given. An experimental test of Ref. [10] in the qubit case has been reported [16], using nuclear magnetic resonance technique.

The geometric phase of entangled states should be another issue worthy of attention. It may hold a potential application in holonomic quantum computation since the study of entangled spin systems effectively allows us to contemplate the design of a solid state quantum computer [17]. Moreover, it was found that one could in principle devise more robust fault-tolerant quantum computations using the notion of geometric phase in designing a conditional phase shift gate [18, 19, 20]. Since the geometric phase depends solely on the geometry of the intrinsic spin space, it is deemed to be less susceptible to noise from the environment. As for the geometric phase of entangled states itself, Sjöqvist [21] considered the geometric phase for a pair of entangled spin- $\frac{1}{2}$ systems in a time-independent uniform magnetic field, and the relative phase for polarization-entangled two-photon systems has been considered by Hessmo and Sjöqvist [22]. Tong *et al.* [23] calculated the geometric phase for a pair of entangled spin- $\frac{1}{2}$ systems in a rotating

* Electronic address: eriks@kvac.uu.se

† Electronic address: phyohch@nus.edu.sg

‡ Electronic address: mericssn@uiuc.edu

magnetic field. Another interesting question, which has been mentioned in the previous papers but has not been completely discussed, is the relation between geometric phase of the entangled state and those of the subsystems.

In this paper, we consider a general entangled bi-partite system under an arbitrary bi-local unitary evolution. We extend the conclusions obtained from the special mode in [23] to the general case. That is, we prove that the geometric phase for cyclic entangled states with non-degenerate eigenvalues under bi-local unitary evolution can always be decomposed into a sum of weighted non-modular pure state phases pertaining to the separable components of the Schmidt decomposition, irrespective of forms of local evolution operators. However, we also see that this property does not manifest itself in the case of non-cyclic evolutions. Moreover, we investigate the relation between the geometric phase of pure entangled states of the system and that of mixed states of the subsystem and conclude that the geometric phase of mixed states of one subsystem is different from that of entangled states in general even if we do not act on the other subsystem. We also point out that the two phases are equal when the evolution operator only acts on the considered subsystem and satisfies conditions where each pure state component of the Schmidt decomposition is parallel transported.

II. NON-CYCLIC GEOMETRIC PHASE OF ENTANGLED STATE

We begin by considering a quantum system S consisting of two subsystems S_a and S_b subject to the bi-local unitary evolution $U(t) = U_a(t) \otimes U_b(t)$. The states of the system belong to the Hilbert space $\mathcal{H} = \mathcal{H}_a \otimes \mathcal{H}_b$, where $\mathcal{H}_a$ and $\mathcal{H}_b$ are two complex Hilbert spaces with dimensions N_a and N_b , respectively. Vectors in $\mathcal{H}_a \otimes \mathcal{H}_b$ can be expanded as Schmidt decompositions. Thus, any normalized initial state of the system can be written as

$$|\Psi(0)\rangle = \sum_{i=1}^N \sqrt{p_i} |\mu_i\rangle \otimes |\nu_i\rangle, \quad (1)$$

where $|\mu_i\rangle$ and $|\nu_i\rangle$ are orthonormal bases of $\mathcal{H}_a$ and $\mathcal{H}_b$, respectively, $N = \min\{N_a, N_b\}$, and the Schmidt coefficients $\{p_i\}$ fulfill $p_1 + \dots + p_N = 1$. When $U_a(t) \otimes U_b(t)$ acts on $|\Psi(0)\rangle$, we obtain

$$|\Psi(t)\rangle = \sum_{i=1}^N \sqrt{p_i} |\mu'_i(t)\rangle \otimes |\nu'_i(t)\rangle, \quad (2)$$

where $|\mu'_i(t)\rangle = U_a(t)|\mu_i\rangle$ and $|\nu'_i(t)\rangle = U_b(t)|\nu_i\rangle$.

Since $|\Psi(t)\rangle$ is a pure state, its geometric phase can be obtained by removing the dynamical phase from the total phase. As we know, when a pure state evolves from $t = 0$ to $t = \tau$ along a path C in projective Hilbert space, the non-adiabatic geometric phase can be obtained as $\gamma(\tau) = \alpha(\tau) - \beta(\tau)$ with total phase $\alpha(\tau) = \arg\langle\Psi(0)|\Psi(\tau)\rangle$ and dynamical phase $\beta(\tau) = -i \int_0^\tau \langle\Psi(t)|\dot{\Psi}(t)\rangle dt$. Hereafter, we use α , β , and γ to mark total, dynamical, and geometric phases, respectively; and we use t , τ , and T to represent instantaneous time, finite time, and period of cyclic evolution, respectively. With Eqs. (1) and (2), we obtain the total, dynamical, and geometric phase of the entangled state $|\Psi(t)\rangle$ under a bi-local unitary evolution $U_a(t) \otimes U_b(t)$ as

$$\alpha_{ab}(\tau) = \arg \sum_{i,j=1}^N \sqrt{p_i p_j} \langle \mu_i | U_a(\tau) | \mu_j \rangle \langle \nu_i | U_b(\tau) | \nu_j \rangle, \quad (3)$$

$$\begin{aligned} \beta_{ab}(\tau) = & \sum_{i=1}^N p_i \left(-i \int_0^\tau \langle \mu_i | U_a^\dagger(t) \dot{U}_a(t) | \mu_i \rangle dt \right) \\ & + \sum_{i=1}^N p_i \left(-i \int_0^\tau \langle \nu_i | U_b^\dagger(t) \dot{U}_b(t) | \nu_i \rangle dt \right), \end{aligned} \quad (4)$$

$$\gamma_{ab}(\tau) = \alpha_{ab}(\tau) - \beta_{ab}(\tau). \quad (5)$$

Eq. (4) entails that the dynamical phase can always be separated into two parts corresponding to the evolution of each of the subsystems S_a and S_b . However, the total phase as well as the geometric phase cannot be separated into two parts in general. The latter observation arises primarily from the entanglement of the two subsystems.

III. CYCLIC GEOMETRIC PHASE OF ENTANGLED STATES

In this section, we specialize the above discussion to cyclic states with non-degenerate Schmidt coefficients p_i . Such states are characterized by the existence of a period T such that

$$|\Psi(T)\rangle = e^{i\alpha_{ab}(T)}|\Psi(0)\rangle, \quad (6)$$

that is,

$$\sum_{i=1}^N \sqrt{p_i} |\mu'_i(T)\rangle \otimes |\nu'_i(T)\rangle = e^{i\alpha_{ab}(T)} \sum_{i=1}^N \sqrt{p_i} |\mu_i\rangle \otimes |\nu_i\rangle. \quad (7)$$

As $U_a(T)$ and $U_b(T)$ are unitary, the vectors $|\mu'_i(T)\rangle$ and $|\nu'_i(T)\rangle$ are also orthonormal bases of $\mathcal{H}_a$ and $\mathcal{H}_b$, respectively. Moreover, under consideration that the p_i 's are non-degenerate, i.e., that $p_i \neq p_j$ for all pairs $i, j \neq i$, the Schmidt decomposition is unique. So, we have $|\mu'_i(T)\rangle \otimes |\nu'_i(T)\rangle = e^{i\alpha_{ab}(T)} |\mu_i\rangle \otimes |\nu_i\rangle$, which implies

$$\alpha_{ab}(T) = \arg [\langle \mu_i | U_a(T) | \mu_i \rangle \langle \nu_i | U_b(T) | \nu_i \rangle], \quad (8)$$

where $\arg \in [0, 2\pi)$. Since the left-hand side of Eq. (8) is independent of the summation index i , $\alpha_{ab}(T)$ can be written as

$$\begin{aligned} \alpha_{ab}(T) &= \sum_{i=1}^N p_i \arg [\langle \mu_i | U_a(T) | \mu_i \rangle \langle \nu_i | U_b(T) | \nu_i \rangle] \\ &= \sum_{i=1}^N p_i [\arg \langle \mu_i | U_a(T) | \mu_i \rangle + \arg \langle \nu_i | U_b(T) | \nu_i \rangle + 2\pi \tilde{n}_i] \end{aligned} \quad (9)$$

where $\tilde{n}_i$ are integers chosen as

$$\begin{aligned} \tilde{n}_i &= 0, \quad \text{if } \arg \langle \mu_i | U_a(T) | \mu_i \rangle + \arg \langle \nu_i | U_b(T) | \nu_i \rangle < 2\pi, \\ \tilde{n}_i &= -1, \quad \text{if } \arg \langle \mu_i | U_a(T) | \mu_i \rangle + \arg \langle \nu_i | U_b(T) | \nu_i \rangle \geq 2\pi. \end{aligned} \quad (10)$$

Substituting Eqs. (4) and (9) into Eq. (5) yields

$$\gamma_{ab}(T) = \sum_{i=1}^N p_i [\gamma_{ai}(T) + \gamma_{bi}(T) + 2\pi n_i], \quad (11)$$

where

$$\begin{aligned} \gamma_{ai}(T) &= \arg \left(\langle \mu_i | U_a(T) | \mu_i \rangle e^{-\int_0^T \langle \mu_i | U_a^\dagger(t) \dot{U}_a(t) | \mu_i \rangle dt} \right), \\ \gamma_{bi}(T) &= \arg \left(\langle \nu_i | U_b(T) | \nu_i \rangle e^{-\int_0^T \langle \nu_i | U_b^\dagger(t) \dot{U}_b(t) | \nu_i \rangle dt} \right), \end{aligned} \quad (12)$$

are just the 2π -modular geometric phases of the pure states $|\mu_i\rangle$ and $|\nu_i\rangle$, respectively. The “winding numbers” n_i are integers and originate from the non-modular nature of the pure state dynamical phases [6] and the integers $\tilde{n}_i$. Eq. (11) shows that the cyclic geometric phase for non-degenerate entangled states under a bi-local unitary evolution can always be decomposed into a sum of weighted pure state phases pertaining to the evolution of each Schmidt component. This is primarily a result of the uniqueness of the Schmidt decomposition, which entails that the Schmidt basis of the initial and final state must be identical if all p_i are different.

The contribution from the non-modular nature of the dynamical phases to the winding numbers n_i can be determined in a history-dependent manner [24, 25] by continuously monitoring the dynamical phase for each separate component of the Schmidt decomposition on the time interval $[0, T]$. In such a procedure, each $2\pi n_i$ is the modulus 2π remainder of the corresponding dynamical phase.

To illustrate the significance of the winding numbers, consider a pair of qubits (two-level systems) being initially in the entangled state

$$|\Psi(0)\rangle = \sqrt{p_0} |0_a\rangle \otimes |0_b\rangle + \sqrt{p_1} |1_a\rangle \otimes |1_b\rangle \quad (13)$$

with $p_0 \neq p_1$, and evolving under influence of the time-independent Hamiltonian

$$H = \hbar\omega|1_a\rangle\langle 1_a| \otimes I_b \quad (14)$$

with $\omega > 0$ and I_b the identity operator on $\mathcal{H}_b$. As the Schmidt components $|0_a\rangle \otimes |0_b\rangle$ and $|1_a\rangle \otimes |1_b\rangle$ are eigenstates of H , their corresponding 2π -modular geometric phases $\{\gamma_{a0}, \gamma_{b0}\}$ and $\{\gamma_{a1}, \gamma_{b1}\}$ all vanish. However, they acquire different dynamical phases causing a non-trivial evolution of the entangled state. For one cycle $T = 2\pi/\omega$ one obtains $n_0 = 0$ and $n_1 = 1$, yielding the total geometric phase

$$\gamma_{ab}(T) = 2\pi p_1, \quad (15)$$

which is non-trivial for entangled states.

IV. COMPARING THE PHASE OF ENTANGLED STATES WITH THAT OF MIXED STATES

The above result concerning the phase of cyclic entangled states may provide some insight regarding the geometric phase of mixed states. Certainly, the state given by Eq. (2) is a pure state with density matrix $\rho_{ab}(t) = |\Psi(t)\rangle\langle\Psi(t)|$. However, if we trace out the state of subsystem S_b , we obtain the reduced density matrix ρ_a corresponding to mixed states of subsystem S_a . By the reduced density matrix, we can deduce the geometric phase $\gamma_a^M(T)$ of a mixed state. Here, we wish to examine, in the case where $U_b(t)$ is the identity map I_b , the relation between $\gamma_{ab}(T)$ and $\gamma_a^M(T)$.

From Eqs. (1) and (2), tracing out the state of subsystem S_b , we obtain the evolution of the reduced density matrix for subsystem S_a as

$$\rho_a(t) = U_a(t)\rho_a(0)U_a^\dagger(t) \quad (16)$$

with

$$\rho_a(0) = \sum_{i=1}^N p_i |\mu_i\rangle\langle\mu_i|. \quad (17)$$

For non-degenerate p_i , the geometric phase of $\rho_a(t)$ is found as [10, 13]

$$\gamma_a^M(\tau) = \arg \left(\sum_{i=1}^N p_i \langle\mu_i|U_a(\tau)|\mu_i\rangle e^{-\int_0^\tau \langle\mu_i|U_a^\dagger(t)\dot{U}_a(t)|\mu_i\rangle dt} \right). \quad (18)$$

This expression is valid for both cyclic and non-cyclic states. In the cyclic case $\tau = T$ we have $|\langle\mu_i|U_a(T)|\mu_i\rangle| = 1$, and the above equation can be written as

$$\gamma_a^M(T) = \arg \left(\sum_{i=1}^N p_i e^{i\gamma_{ai}(T)} \right), \quad (19)$$

where $\gamma_{ai}(T)$ is the geometric phase of the pure state $|\mu_i\rangle$. Comparing Eq. (19) with Eq. (11) for $U_b(t) = I_b$, we find $\gamma_a^M(T) \neq \gamma_{ab}(T)$. Thus, the cyclic geometric phase of the whole system is in general different from that of the mixed state of the considered subsystem, basically because the former is a weighted sum of pure state phases while the latter is a weighted sum of pure state phase factors.

Similarly, for non-cyclic evolution with $U(t) = U_a(t) \otimes I_b$, we have in general

$$\gamma_{ab}(\tau) \equiv \gamma_a(\tau) = \arg \sum_{i=1}^N p_i \langle\mu_i|U_a(\tau)|\mu_i\rangle + i \int_0^\tau \sum_{i=1}^N p_i \langle\mu_i|U_a^\dagger(t)\dot{U}_a(t)|\mu_i\rangle dt \quad (20)$$

and

$$\gamma_a^M(\tau) = \arg \left(\sum_{i=1}^N p_i |\langle\mu_i|U_a(\tau)|\mu_i\rangle| e^{i\gamma_{ai}(\tau)} \right). \quad (21)$$

Thus, the geometric phase of the system S is dependent upon $|\mu_i\rangle$ of subsystem S_a but independent of $|\nu_i\rangle$ of S_b . Only the evolution of subsystem S_a contributes to the geometric phase of the pure state system. Yet, we find that, even in the case when the system's geometric phase is completely determined by the evolution of subsystem S_a and the Schmidt coefficients $\{p_i\}$, $\gamma_a(\tau)$ is generally different from $\gamma_a^M(\tau)$ both in the cyclic and non-cyclic case. It may seem unexpected, because only subsystem S_a experiences a unitary evolution while subsystem S_b is unaffected. The geometric phase of the system S is attributed to $U_a(t)$ only, and it seems natural to expect the phase obtained by the system to be same as that obtained by subsystem S_a , while regarding it as mixed state. However, they are different. This shows that the geometric phase of an entangled bi-partite system is always affected by both subsystems.

V. PHASE RELATIONS UNDER PARALLEL TRANSPORT CONDITIONS

In this section, we again restrict our discussion to the case where $U(t) = U_a(t) \otimes I_b$. As pointed out above, even in this case, the geometric phase of the system, which is determined only by the evolution of subsystem S_a and the Schmidt coefficients $\{p_i\}$, is generally different from that of the corresponding mixed state. We now try to find the reason for the difference and give conditions under which the two phases are equal.

Under the evolution $U(t) = U_a(t) \otimes I_b$, the state of the whole system is $|\Psi(t)\rangle = \sum_{i=1}^N \sqrt{p_i} [U_a(t)|\mu_i\rangle] \otimes |\nu_i\rangle$, which can also be expressed as the density matrix

$$\rho(t) = \sum_{i=1}^N \sqrt{p_i p_j} [U_a(t)|\mu_i\rangle \langle \mu_j| U_a^\dagger(t)] \otimes |\nu_i\rangle \langle \nu_j|. \quad (22)$$

The corresponding mixed state of the subsystem S_a is

$$\rho_a(t) = \text{Tr}_b \rho(t) = \sum_{i=1}^N p_i U_a(t) |\mu_i\rangle \langle \mu_i| U_a^\dagger(t). \quad (23)$$

When $U_a(t)$ is given, the states $\rho(t)$ and $\rho_a(t)$ are definite, but when $\rho(t)$ or $\rho_a(t)$ is given, the evolution operator is not unique. That is, for a given path in state space, there are infinitely many unitary operators that realize the same path and so give the same geometric phase. All the operators form an equivalence set: two evolution operators are ‘equivalent’ if and only if they realize the same path. If we know any one operator out of the equivalence set, say $\tilde{U}_a(t)$, we can write down all the operators of the set. For $\rho(t)$, the equivalence set is

$$\mathcal{S}_1 = \left\{ \tilde{U}_a(t) e^{i\theta(t)} \right\}, \quad (24)$$

where $\theta(t)$ is an arbitrary real-valued gauge function of t with $\theta(0) = 0$. For state $\rho_a(t)$, the equivalence set is

$$\mathcal{S}_2 = \left\{ \tilde{U}_a(t) \sum_{i=1}^N e^{i\theta_i(t)} |\mu_i\rangle \langle \mu_i| \right\}, \quad (25)$$

where $\theta_i(t)$, $i = 1, 2, \dots, N$, are arbitrary real-valued gauge functions of t with $\theta_i(0) = 0$. We see that the two sets are different in general and $\mathcal{S}_1 \subset \mathcal{S}_2$, which shows that the evolution operators that give the same path for $\rho_a(t)$ may give different paths for $\rho(t)$. So the two kinds of geometric phases $\gamma_a(\tau)$ and $\gamma_a^M(\tau)$ cannot be the same in general, otherwise they should be associated with the same equivalence sets of evolution operators. To see exactly the difference between the two phases, we substitute

$$U_a(t) = \tilde{U}_a(t) \sum_{i=1}^N e^{i\theta_i(t)} |\mu_i\rangle \langle \mu_i| \quad (26)$$

into Eqs. (18) and (20), respectively, and get

$$\gamma_a^M(\tau) = \arg \left(\sum_{i=1}^N p_i \langle \mu_i | \tilde{U}_a(\tau) | \mu_i \rangle e^{-\int_0^\tau \langle \mu_i | \dot{\tilde{U}}_a^\dagger(t) \dot{\tilde{U}}_a(t) | \mu_i \rangle dt} \right), \quad (27)$$

$$\begin{aligned} \gamma_a(\tau) &= \arg \sum_{i=1}^N p_i \langle \mu_i | \tilde{U}_a(\tau) | \mu_i \rangle e^{i\theta_i(\tau)} \\ &\quad + i \int_0^\tau \sum_{i=1}^N p_i \langle \mu_i | \dot{\tilde{U}}_a^\dagger(t) \dot{\tilde{U}}_a(t) | \mu_i \rangle dt - \sum_{i=1}^N p_i \theta_i(\tau). \end{aligned} \quad (28)$$

We see that $\gamma_a^M(\tau)$ is invariant under choice of member in $\mathcal{S}_2$, while $\gamma_a(\tau)$ is not as it depends upon $\theta_i(t)$.

With the above analysis, we can conclude that, for a given local evolution operator $U_a(t)$, the two phases $\gamma_a(\tau)$ and $\gamma_a^M(\tau)$ are different in general. But when can they be the same, that is, for what kinds of $U_a(t)$ can we consider the two phases to be the same? We prove that when the evolution operator $U_a(t)$ satisfies ‘the stronger parallel transport conditions’ [10]

$$\langle \mu_i | U_a^\dagger(t) \dot{U}_a(t) | \mu_i \rangle = 0, \quad i = 1, 2, \dots, N, \quad (29)$$

the two phases are the same. Substituting Eq. (29) into Eqs. (18) and (20), we find

$$\gamma_a(\tau) = \gamma_a^M(\tau) = \arg \left(\sum_{i=1}^N p_i \langle \mu_i | U_a(\tau) | \mu_i \rangle \right), \quad (30)$$

and

$$\gamma_b(\tau) = \gamma_b^M(\tau) = 0. \quad (31)$$

Eqs. (30) and (31) show that the two kinds of geometric phase for unitarities of the form $U_a(t) \otimes I_b$ are always the same when the evolution operator $U_a(t)$ satisfies the stronger parallel transport conditions Eq. (29).

Substituting $U_a(t) = V_a(t) \sum_{i=1}^N e^{i\theta_i(t)} |\mu_i\rangle\langle\mu_i|$ into Eq. (29), we get the general form of evolution operators satisfying the stronger parallel transport conditions

$$U_a(t) = V_a(t) \sum_{i=1}^N |\mu_i\rangle\langle\mu_i| e^{-\int_0^t \langle \mu_i | V_a^\dagger(t) \dot{V}_a(t) | \mu_i \rangle dt}, \quad (32)$$

where $V_a(t)$ is an arbitrary unitary operator. So we see that when $U_a(t)$ holds the form of Eq. (32), the geometric phase of the pure state of the system under the evolution $U(t) = U_a(t) \otimes I_b$ is the same as that of the mixed state of subsystem S_a under the evolution $U_a(t)$. The phase relations are shown as Eqs. (30) and (31).




FIG. 1: Two-particle interferometry setups to measure phases for internal degrees of freedom in entangled pure state $|\Psi(0)\rangle$. In the upper panel, the pure state phase $\arg\langle\Psi(0)|U_a(\tau)\otimes I_b|\Psi(0)\rangle$ is measured as a shift in the coincidence interference oscillations obtained at detectors D_a and D_b by varying the $U(1)$ phase χ . In the lower panel, the mixed state phase $\arg\text{Tr}(\rho_a(0)U_a(\tau))$, with $\rho_a(0) = \text{Tr}_b|\Psi(0)\rangle\langle\Psi(0)|$ the mixed input state of subsystem a , is similarly measured as a shift at D_a by ignoring the subsystem b .

It might also be useful to consider the problem from an operational point of view. Let us first consider the two-particle Franson type interferometry setup shown in the upper panel of Fig. 1, with internal pure state $|\Psi(0)\rangle$ affected

by $U_a(t) \otimes I_b$ in the longer arms and by a $U(1)$ shift χ to one of the shorter arms. To observe the relative phase between $|\Psi(0)\rangle$ and $|\Psi(\tau)\rangle = U_a(\tau) \otimes I_b |\Psi(0)\rangle$, we require that the source produces particle pairs randomly. Then, if the particles arrive at the two detectors D_a and D_b simultaneously, they both either took the upper path or the lower path, so that the intensity detected in coincidence at D_a and D_b becomes [22]

$$\begin{aligned} \mathcal{I}_{ab} &\propto \left| e^{i\chi} |\Psi(0)\rangle + |\Psi(\tau)\rangle \right|^2 \\ &= 2 + 2 \left| \langle \Psi(0) | U_a(\tau) \otimes I_b | \Psi(0) \rangle \right| \cos \left[\chi - \arg \langle \Psi(0) | U_a(\tau) \otimes I_b | \Psi(0) \rangle \right]. \end{aligned} \quad (33)$$

Thus, the relative phase $\arg \langle \Psi(0) | U_a(\tau) \otimes I_b | \Psi(0) \rangle$ shifts the interference oscillations when χ varies. To observe the phase of $\rho_a(t) = \text{Tr}_b |\Psi(t)\rangle \langle \Psi(t)|$ acquired by subsystem a alone, we instead use the setup shown in the lower panel of Fig. 1 with the same source but where subsystem b is ignored. Here, the output intensity at D_a reads [10]

$$\mathcal{I}_a \propto 2 + 2 \left| \text{Tr}(\rho_a(0) U_a(\tau)) \right| \cos \left[\chi - \arg \text{Tr}(\rho_a(0) U_a(\tau)) \right]. \quad (34)$$

Note that $\text{Tr}(\rho_a(0) U_a(\tau)) = \langle \Psi(0) | U_a(\tau) \otimes I_b | \Psi(0) \rangle$ due to the triviality of the evolution of the b subsystem.

Suppose now we first arrange these experiments so that $U_a(t)$ yields the pure state parallel transport condition $\langle \Psi(0) | U_a^\dagger(t) \dot{U}_a(t) \otimes I_b | \Psi(0) \rangle = 0$. In this case, the interference pattern obtained by measuring on both subsystems in coincidence would be shifted by the pure state geometric phase $\gamma_a(\tau)$. On other hand, measuring only on subsystem a would yield the same shift, but the interpretation is different: it is the total phase $\alpha_a^M(\tau)$ acquired by $\rho_a(\tau)$, which is in general at variance with $\gamma_a^M(\tau)$ as this latter phase is based upon the stronger parallel transport conditions. The situation is different when setting up $U_a(t)$ so as to parallel transport $\rho_a(t)$, i.e., by implementing $U_a(t)$ according to Eq. (32). Here, the coincidence and marginal interference pattern are shifted by $\gamma_a(\tau)$ and $\gamma_a^M(\tau)$, respectively, in accordance with the above analysis.

VI. CONCLUSION AND REMARKS

We have discussed geometric phases of entangled states of bi-partite systems under bi-local unitary evolution and of the mixed states of their subsystems. We conclude:

1. The cyclic geometric phase for entangled states with non-degenerate eigenvalues under bi-local unitary evolution can always be decomposed into a sum of weighted non-modular pure state phases pertaining to the separable components of the Schmidt decomposition, irrespective of forms of local evolution operators, though the same cannot be said for the non-cyclic geometric phase.
2. The mixed state geometric phase of one subsystem is generally different from that of the entangled state even by keeping the other subsystem fixed, though it seems as if the two phases might be same. However, when the evolution operator satisfies the stronger parallel transport conditions for mixed states, the two phases are the same, and the general form of the operators are given.

The difference between geometric phases of bi-partite systems and their parts has its primary cause in entanglement and thus vanishes in the limit of separable states. We hope that the present analysis may trigger multi-particle experiments to test the difference between phases of entangled systems and their subsystems.

Acknowledgments

The work by Tong was supported by NUS Research Grant No. R-144-000-054-112. This work is also supported in part by the Agency for Science, Technology and Research, Singapore under the ASTAR Grant No. 012-104-0040 (WBS: R-144-000-071-305). E.S. acknowledges financial support from the Swedish Research Council. M.E. acknowledge financial support from the Foundation BLANCEFLOR Boncompagni-Ludovisi, née Bildt.

-
- [1] M.V. Berry, Proc. R. Soc. London Ser. A **392**, 45 (1984).
[2] B. Simon, Phys. Rev. Lett. **51**, 2167 (1983).

- [3] Y. Aharonov and J. Anandan, Phys. Rev. Lett. **58**, 1593 (1987).
- [4] J. Anandan and Y. Aharonov, Phys. Rev. D **38**, 1863 (1988).
- [5] J. Samuel and R. Bhandari, Phys. Rev. Lett. **60**, 2339 (1988).
- [6] N. Mukunda and R. Simon, Ann. Phys. (N.Y.) **228**, 205 (1993).
- [7] A.K. Pati, Phys. Lett. A **202**, 40 (1995).
- [8] N. Manini and F. Pistolesi, Phys. Rev. Lett. **85**, 3067 (2000).
- [9] A. Uhlmann, Rep. Math. Phys. **24**, 229 (1986); Lett. Math. Phys. **21**, 229 (1991).
- [10] E. Sjöqvist, A.K. Pati, A. Ekert, J.S. Anandan, M. Ericsson, D.K.L. Oi, and V. Vedral, Phys. Rev. Lett. **85**, 2845 (2000).
- [11] P.B. Slater, Lett. Math. Phys. **60**, 123 (2002).
- [12] M. Ericsson, A.K. Pati, E. Sjöqvist, J. Brännlund, and D.K.L. Oi, Phys. Rev. Lett. **91**, 090405 (2003).
- [13] K. Singh, D.M. Tong, K. Basu, J.L. Chen, and J.F. Du, Phys. Rev. A **67**, 032106 (2003).
- [14] S. Filipp and E. Sjöqvist, Phys. Rev. Lett. **90**, 050403 (2003).
- [15] M. Ericsson, E. Sjöqvist, J. Brännlund, D.K.L. Oi, and A.K. Pati, Phys. Rev. A **67**, 020101(R) (2003).
- [16] J.F. Du, P. Zou, M. Shi, L.C. Kwek, J.-W. Pan, C.H. Oh, A. Ekert, D.K.L. Oi, and M. Ericsson, Phys. Rev. Lett. **91**, 100403 (2003).
- [17] G. Burkard and D. Loss, Phys. Rev. Lett. **88**, 047903 (2002); D.P. DiVincenzo, G. Burkard, D. Loss, and E.V. Sukhorukov, in *Quantum Mesoscopic Phenomena and Mesoscopic Devices in Microelectronics*, edited by I.O. Kulik and R. Ellialtıoglu (NATO Advanced Study Institute, Turkey, 1999).
- [18] J.A. Jones, V. Vedral, A. Ekert, and G. Castagnoli, Nature (London) **403**, 869 (2000).
- [19] A.K. Ekert, M. Ericsson, P. Hayden, H. Inamori, J.A. Jones, D.K.L. Oi, and V. Vedral, J. Mod. Opt. **47**, 2501 (2000).
- [20] X.B. Wang and K. Matsumoto, Phys. Rev. Lett. **87**, 097901 (2001).
- [21] E. Sjöqvist, Phys. Rev. A **62**, 022109 (2000).
- [22] B. Hessmo and E. Sjöqvist, Phys. Rev. A **62**, 062301 (2000).
- [23] D.M. Tong, L.C. Kwek, and C.H. Oh, J. Phys. A **36**, 1149 (2003).
- [24] R. Bhandari, Phys. Lett. A **157**, 221 (1991); Phys. Lett. A **171**, 262 (1992); Phys. Lett. A **171**, 267 (1992); Phys. Lett. A **180**, 15 (1993).
- [25] R. Bhandari, Phys. Rev. Lett. **89**, 268901 (2002); J. Anandan, E. Sjöqvist, A.K. Pati, A. Ekert, M. Ericsson, D.K.L. Oi, and V. Vedral, Phys. Rev. Lett. **89**, 268902 (2002).